



SACLAY



DIRECTION DES SCIENCES DE LA MATIERE,
DEPARTEMENT DE RECHERCHE SUR L'ETAT CONDENSE,
LES ATOMES ET LES MOLECULES,
SERVICE DE PHYSIQUE ET DE CHIMIE DES SURFACES ET DES INTERFACES

SEMINAIRE *

Vendredi 29 juin 2007 à 11h00

Bâtiment 466, salle 111 - CEA Saclay, 91191, Gif sur Yvette

Anticracks in the venation network of leaves?

P. M. Reis

PMMH, ESPCI

Invité par D. BONAMY

Résumé:

Thus far, existing models of venation in leaves are entirely biochemical, involving hormonal diffusive processes. These are, however, unable to capture some crucial structural features of the network of vascular bundles such as the existence of reconnection loops. Couder et. al. [1] have recently highlighted the striking similarities between leaf venation and fracture patterns in drying gels, suggesting that a tensorial mechanism may be at play. It is known that, in the initial stages of the formation of vascular bundles, the promesophyll ('bulk' of the leaf) is under compression since it grows at a faster rate than the protoderm ('skin' of the leaf). Hence, to take this analogy further, we introduce the concept of an 'anticrack': a localization of deformation under compressive stresses. We have developed an experimental system to develop and explore this concept as when a solid foam is compressed, either uniaxially or biaxially. We analyze the resulting anticrack networks and relate them to the fracture and venation counterparts. For this purpose we use a high resolution image correlation technique to measure the statistics of the localization zones, structural hierarchy and reconnection loops.

*** SERA PRECEDE D'UNE PAUSE-CAFE A PARTIR DE 10H30**